

## DENTAL - EXPOSED PULP

(ISS MED/3A - ALL/FIN)

Page 1 of 2 pages

I

1. Anesthetize tooth.  
Refer to {DENTAL - INJECTION TECHNIQUE} (SODF: ISS MED: DENTAL).
2. Unstow:  
AMP  
(blue)      Toothache Kit (Dental-7)  
                 Tubex Injector (Injectables-14)  
                 Gauze Pad (P3-B4)
3. From Toothache Kit, unstow:  
                 Eugenol Anesthetic Medication Cartridge (1)  
                 Tweezers  
                 Cotton Pellet (1)

### NOTE

Eugenol Anesthetic Drops is an eye irritant (Tox Level 1). Use sparingly.

4. Assemble Tubex Injector with Eugenol Anesthetic Medication Cartridge.  
Refer to {INJECTIONS - TUBEX INJECTOR} (SODF: ISS MED: INJECTIONS/IV).
5. Moisten Cotton Pellet with Eugenol Anesthetic by ejecting a small amount of medication from cartridge, keeping tip in direct contact of Cotton Pellet at all times.
6. Recap Tubex if useable medication remains in cartridge.
7. Squeeze excess Eugenol Anesthetic out of Cotton Pellet with Gauze Pad.
8. Place moistened Cotton Pellet on exposed pulp.  
Do not cover with anything else.

### WARNING

Do not place Temporary Filling over exposed pulp.

9. Successive placement of Eugenol Anesthetic should not require additional local anesthesia.

**DENTAL - EXPOSED PULP**

(ISS MED/3A - ALL/FIN)

Page 2 of 2 pages

10. Avoid tooth when chewing and keep food out of cavity.
11. Restow Toothache Kit contents.
12. Contact Surgeon for further instructions.