

ABRASIONS

(ISS MED/3A - ALL/FIN)

Page 1 of 2 pages

- AMP (blue) 1. Unstow:
Gauze Pads (P3-B4) (Surgical Supply-4)
Polysporin Ointment (P2-A12)
Erythromycin Ointment
Solutio Viride Nitens
Ethyl Alcohol
Bandaid (P4-B3)
Kling Dressing (Surgical Supply-6) (P3-B5)
- AMP (blue) 2. Clean affected area thoroughly with soap and water, followed by Ethyl Alcohol.
Blot dry gently with Gauze Pads.
3. Photodocument the wound.
4. As advised from ground, apply Solutio Viride Nitens to skin around wound.
5. Dress wound with Polysporin or Erythromycin Ointment.
6. Cover with Bandaid or with Gauze Pad for large abrasions.
7. Use Tape or wrap with Kling Dressing to hold in place.
8. Repeat steps 1 --- 3.
Inspect abrasion once daily or when abrasion becomes wet or contaminated.
9. Notify Surgeon at next routine PMC.
10. Continue to photodocument every 24 hours.
Archive and downlink.
11. For slow-healing wounds, use Solcoseryl Ointment.

ABRASIONS

(ISS MED/3A - ALL/FIN)

Page 2 of 2 pages

Medical Kit Solcoseryl Ointment - Used to accelerate regenerative processes.
with
Ointments

Dose: Apply to wound area 1-2 times/day

Possible side effects

Local mild skin irritation, burning