

BEHAVIORAL - ACUTE PSYCHOSIS

(ISS MED/3A - ALL/FIN)

Page 1 of 3 pages

I

NOTE

Contact Surgeon before giving any medication marked with an asterisk. In an emergency or during Loss of Signal, begin appropriate treatment; then call Surgeon as soon as possible.

- ALSP (red)
1. Unstow:
 - Drug Subpack
 - Gray Tape
 - Bungees
 - Towels
 2. Talk with the patient while you are restraining him.
Explain what you are doing, and that you are using a restraint to ensure that he is safe.
Restrain patient using Gray Tape around wrists, ankles, and use a bungee around the torso.
If necessary to restrain the head, place a rolled towel under the neck and restrain with Gray Tape.
- AMP (blue)
3. Administer 10 mg Haldol (Haloperidol) Oral (P4-B6) - Potent tranquilizer
If patient is uncooperative with taking oral dose of Haldol (Haloperidol), give Haldol (Haloperidol) IM as noted in step 4.

Possible side effects
Low blood pressure, rapid heart rate, headache, involuntary movements; increased body temperature
- ALSP (red)
4. Administer 5 mg/ml of Haldol (Haloperidol) Injectables IM (Drug 27,28). Refer to {INJECTIONS - INTRAMUSCULAR} (SODF: ISS MED: INJECTIONS/IV).

NOTE

Backup CMO should remain with restrained crewmember at all times. Confirm presence of radial and pedal pulses. If pulses are weak or faint, loosen restraints.

BEHAVIORAL - ACUTE PSYCHOSIS

(ISS MED/3A - ALL/FIN)

Page 2 of 3 pages



Figure 1.- Intramuscular Injection.

1. Prepare injection site with Alcohol Pad.
2. Stabilize hand on arm and enter skin gently but rapidly perpendicular to surface, about 2-3 cm deep.
3. Withdraw the plunger slightly to look for blood return to ensure that the needle is not in a vessel. If so, withdraw and try again at a diferent site.
4. Inject Medication Cartridge.

- AMP
(blue)
5. Administer 5 mg Valium (Diazepam) Oral (P1-A12) - Sedative, anticonvulsant, antiseizure drug

If patient is uncooperative with taking oral dose of Valium (Diazepam), give IM Valium (Diazepam) as noted in step 6.

Possible side effects

Drowsiness, fatigue, nausea, slow heart rate, blurred vision.
Romazicon is antidote for overdose.

6. Administer 5 mg/ml (half an ampule) Valium (Diazepam) IM (Drug-8,9,10). Refer to {[INJECTIONS - INTRAMUSCULAR](#)} (SODF: ISS MED: INJECTIONS/IV).

NOTE

Crewmember will be sufficiently sedated. Severe muscle spasms may occur. If present, give 50 mg Bendryl (Diphenhydramine) IM and maintain adequate fluid hydration.

BEHAVIORAL - ACUTE PSYCHOSIS

(ISS MED/3A - ALL/FIN)

Page 3 of 3 pages

Possible side effects of Benadryl

Drowsiness, inability to concentrate, dry mouth, blurred vision, rash, sensitivity to light, headache, rapid heart rate, dizziness, fatigue

7. Monitor and record vital signs while the patient is restrained:

If blood pressure, pulse, or breathing is abnormal or unstable, check vital signs approximately every 5 minutes and record values with time.

If and when blood pressure, pulse, and breathing are stable for two or three readings, decrease frequency to every 15 minutes until advised by ground.

	Time (minutes)						
	Pulse (radial (#), pedal (±))						
ALSP	Blood Presssure (ALSP-4)						
	Respiratory Rate						
AMP	Temperature (Skin Temp Monitors, P3-A7)						

8. Contact Surgeon to discuss crewmember status and restraint removal.