

SLEEP MEDICATION OVERDOSE - EMERGENCY

(ISS MED/3A - ALL/FIN)

Page 1 of 2 pages

I

SIGNS

Confusion
Lethargy
Difficulty waking

SYMPTOMS

Decreased respiratory rate
Low blood pressure
Slow heart rate

1. If not breathing, go to {CARDIOPULMONARY RESUSCITATION: CPR - ORAL AIRWAY INSERTION - EMERGENCY} (SODF: ISS MED: EMERGENCY).
2. Perform vital signs and record as necessary every 5 minutes.

	Time (minutes)	0	5	10	15	20	25	30
ALSP	Blood Pressure (ALSP-4)							
	Pulse							
	Respiratory Rate							
ALSP	Temperature (Assessment-4)							
ALSP	Pulse Oximeter (Assessment-1)							

WARNING

Recurrence of sedation can occur if large amounts of sedatives were ingested. Close monitoring is needed for several hours after use. If sedation recurs, repeat dosing below no more often than every 20 minutes.

3. Romazicon (Flumazenil) Injectable (Drug-19 --- 22, 58) - Antidote for overdose of Restoril or Valium
Refer to {INJECTIONS - BUTTERFLY NEEDLE} (SODF: ISS MED: INJECTIONS/IV).

Dose: Inject 2 ml over 30 seconds.
Wait 30 seconds.

SLEEP MEDICATION OVERDOSE - EMERGENCY

(ISS MED/3A - ALL/FIN)

Page 2 of 2 pages

If no response, repeat with 3 ml over 30 seconds.

Wait 30 seconds.

If no response, repeat 5 ml over 30 seconds.

If still no response, then symptoms are unlikely due to sedative overdose.

Possible side effects

Dizziness, sweating, headache, blurred vision, nausea, vomiting, agitation

3. Contact Surgeon.